



JEE-MAIN – JUNE, 2022

(Held On Tuesday 26th June, 2022)

TIME : 3 : 00 PM to 6 : 00 PM

Mathematics

Test Pattern : JEE-MAIN

Maximum Marks : 120

Topic Covered: FULL SYLLABUS

Important instruction:

1. Use Blue / Black Ball point pen only.
2. There are three sections of equal weightage in the question paper **Physics, Chemistry** and **Mathematics** having 30 questions in each subject. Each paper have 2 sections A and B.
3. You are awarded +4 marks for each correct answer and -1 marks for each incorrect answer.
4. Use of calculator and other electronic devices is not allowed during the exam.
5. No extra sheets will be provided for any kind of work.

Name of the Candidate (in Capitals) _____

Father's Name (in Capitals) _____

Form Number : in figures _____

: in words _____

Centre of Examination (in Capitals): _____

Candidate's Signature: _____ Invigilator's Signature : _____

Rough Space

YOUR TARGET IS TO SECURE GOOD RANK IN JEE-MAIN

Corporate Office : **ALLEN Digital Pvt. Ltd.**, "One Biz Square", A-12 (a), Road No. 1, Indraprastha Industrial Area, Kota (Rajasthan) INDIA-324005

+91-9513736499 | +91-7849901001 | wecare@allendigital.in | www.allendigital.in

FINAL JEE-MAIN EXAMINATION – JUNE, 2022

(Held On Sunday 26th June, 2022)

TIME : 03 : 00 PM to 06 : 00 PM

MATHEMATICS

TEST PAPER WITH SOLUTION

SECTION-A

1. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined as $f(x) = x - 1$ and $g : \mathbb{R} - \{1, -1\} \rightarrow \mathbb{R}$ be defined as $g(x) = \frac{x^2}{x^2 - 1}$.

Then the function fog is :

- (A) one-one but not onto function
- (B) onto but not one-one function
- (C) both one-one and onto function
- (D) neither one-one nor onto function

Official Ans. by NTA (D)

Allen Ans. (D)

Sol. $f(x) = x - 1$; $g(x) = \frac{x^2}{x^2 - 1}$

$$f(g(x)) = g(x) - 1$$

$$= \frac{x^2}{x^2 - 1} - 1 = \frac{x^2 - x^2 + 1}{x^2 - 1}$$

$$f(g(x)) = \frac{1}{x^2 - 1}; x \neq \pm 1, \text{ even function}$$

→ Hence $f(g(x))$ is many one function

$$y = \frac{1}{x^2 - 1}$$

$$y \cdot x^2 - y = 1$$

$$x^2 = \left(\frac{1+y}{y}\right)$$

$$\left(\frac{1+y}{y}\right) \geq 0$$



Range:- $y \in (-\infty, -1] \cup (0, \infty)$

Hence, Range ≠ Co-domain $\Rightarrow f(g(x))$ is into function

2. If the system of equations $\alpha x + y + z = 5$, $x + 2y + 3z = 4$, $x + 3y + 5z = \beta$,

has infinitely many solutions, then the ordered pair (α, β) is equal to :

- (A) (1, -3) (B) (-1, 3)
- (C) (1, 3) (D) (-1, -3)

Official Ans. by NTA (C)

Allen Ans. (C)

For infinitely many solutions,

$$\Delta = 0 = \Delta_x = \Delta_y = \Delta_z$$

Sol. $\Delta = \begin{vmatrix} \alpha & 1 & 1 \\ 1 & 2 & 3 \\ 1 & 3 & 5 \end{vmatrix} = 0$

$$\Rightarrow \alpha(10 - 9) - 1(5 - 3) + 1(3 - 2) = 0$$

$$\Rightarrow \alpha - 2 + 1 = 0$$

$$\Rightarrow \alpha = 1$$

$$\Delta_x = \begin{vmatrix} 5 & 1 & 1 \\ 4 & 2 & 3 \\ \beta & 3 & 5 \end{vmatrix} = 0$$

$$\Rightarrow 5(10 - 9) - 1(20 - 3\beta) + 1(12 - 2\beta) = 0$$

$$\Rightarrow 5 - 20 + 3\beta + 12 - 2\beta = 0$$

$$\Rightarrow -3 + \beta = 0$$

$$\Rightarrow \beta = 3$$

3. If $A = \sum_{n=1}^{\infty} \frac{1}{(3+(-1)^n)^n}$ and $B = \sum_{n=1}^{\infty} \frac{(-1)^n}{(3+(-1)^n)^n}$, then

$\frac{A}{B}$ is equal to :

- (A) $\frac{11}{9}$ (B) 1 (C) $-\frac{11}{9}$ (D) $-\frac{11}{3}$

Official Ans. by NTA (C)

Allen Ans. (C)

Sol. $A = \left(\frac{1}{2} + \frac{1}{4^2} + \frac{1}{2^3} + \frac{1}{4^4} + \dots \right)$

$$A = \left(\frac{1}{2} + \frac{1}{2^3} + \dots \right) + \left(\frac{1}{4^2} + \frac{1}{4^4} + \dots \right)$$

$$A = \left(\frac{\frac{1}{2}}{1 - \frac{1}{4}} + \frac{\frac{1}{16}}{1 - \frac{1}{16}} \right)$$

$$\Rightarrow A = \frac{1}{2} \times \frac{4}{3} + \frac{1}{16} \times \frac{16}{15} \Rightarrow A = \frac{11}{15}$$

$$B = \frac{-1}{2} + \frac{1}{4^2} + \frac{-1}{2^3} + \frac{1}{4^4} + \dots$$

$$B = \left(\frac{-1}{2} + \frac{-1}{2^3} + \dots \right) + \left(\frac{1}{4^2} + \frac{1}{4^4} + \dots \right)$$

$$B = \frac{-\frac{1}{2}}{1 - \frac{1}{4}} + \frac{\frac{1}{16}}{1 - \frac{1}{16}}$$

$$\Rightarrow B = -\frac{1}{2} \times \frac{4}{3} + \frac{1}{16} \times \frac{16}{15}$$

$$B = -\frac{9}{15}$$

$$\frac{A}{B} = \frac{11}{15} \times \frac{15}{(-9)}$$

$$\frac{A}{B} = -\frac{11}{9}$$

4. $\lim_{x \rightarrow 0} \frac{\cos(\sin x) - \cos x}{x^4}$ is equal to :

- (A) $\frac{1}{3}$ (B) $\frac{1}{4}$ (C) $\frac{1}{6}$ (D) $\frac{1}{12}$

Official Ans. by NTA (C)

Allen Ans. (C)

Sol. $\lim_{x \rightarrow 0} \frac{\cos(\sin x) - \cos x}{x^4} : \left(\frac{0}{0} \right)$

$$\lim_{x \rightarrow 0} \left(\frac{2 \cdot \sin\left(\frac{x + \sin x}{2}\right) \cdot \sin\left(\frac{x - \sin x}{2}\right)}{x^4} \right)$$

$$\lim_{x \rightarrow 0} 2 \left(\frac{\sin\left(\frac{x + \sin x}{2}\right)}{\left(\frac{x + \sin x}{2}\right)} \right) \left(\frac{\sin\left(\frac{x - \sin x}{2}\right)}{\left(\frac{x - \sin x}{2}\right)} \right) \left(\frac{\frac{x + \sin x}{2}}{x^4} \right) \left(\frac{\frac{x - \sin x}{2}}{x^4} \right)$$

$$\lim_{x \rightarrow 0} \left(\frac{x^2 - \sin^2 x}{2x^4} \right) : \left(\frac{0}{0} \right)$$

Apply L-Hopital Rule :

$$\lim_{x \rightarrow 0} \frac{2x - 2 \sin x \cos x}{2 \cdot 4 \cdot x^3}$$

$$\lim_{x \rightarrow 0} \frac{2x - \sin 2x}{8x^3} ; 0 : \text{Again apply L-Hopital rule}$$

$$\lim_{x \rightarrow 0} \frac{2 - 2 \cos(2x)}{8(3)x^2}$$

$$\lim_{x \rightarrow 0} \frac{2(1 - \cos(2x))}{24(4x^2)} \times 4 \Rightarrow \frac{2}{24} \times \frac{1}{2} \times 4 \Rightarrow \frac{1}{6}$$


5. Let $f(x) = \min \{1, 1 + x \sin x\}$, $0 \leq x \leq 2\pi$. If m is the number of points, where f is not differentiable and n is the number of points, where f is not continuous, then the ordered pair (m, n) is equal to

- (A) (2, 0) (B) (1, 0)
 (C) (1, 1) (D) (2, 1)

Official Ans. by NTA (B)

Allen Ans. (B)

Sol.



No. of non-differentiable points = 1 (m)

No. of not continuous points = 0 (n)

(m, n) = (1, 0)

6. Consider a cuboid of sides $2x$, $4x$ and $5x$ and a closed hemisphere of radius r . If the sum of their surface areas is a constant k , then the ratio $x : r$, for which the sum of their volumes is maximum, is :

- (A) 2 : 5 (B) 19:45 (C) 3 : 8 (D) 19 : 15

Official Ans. by NTA (B)

Allen Ans. (B)

Sol. Surface area = $76x^2 + 3\pi r^2 = \text{constant } (K)$

$$V = 40x^3 + \frac{2}{3}\pi r^3$$

$$[76x^2 + 3\pi r^2 = K]$$

$$r^2 = \frac{K - 76x^2}{3\pi}$$

$$r = \left(\frac{K - 76x^2}{3\pi} \right)^{\frac{1}{2}}$$

$$V = 40x^3 + \frac{2}{3}\pi \left(\frac{K - 76x^2}{3\pi} \right)^{\frac{3}{2}}$$

$$\frac{dV}{dx} = 120x^2 + \frac{2}{3}\pi \cdot \frac{3}{2} \left(\frac{K - 76x^2}{3\pi} \right)^{\frac{1}{2}} \cdot \left(\frac{-76(2x)}{3\pi} \right)$$

Put

$$\frac{dV}{dx} = 0 \Rightarrow 120x^2 + \frac{2}{3}\pi \cdot \frac{3}{2} \left(\frac{K - 76x^2}{3\pi} \right)^{\frac{1}{2}} \cdot \left(\frac{-76(2x)}{3\pi} \right) = 0$$

$$\Rightarrow 120x^2 = \frac{152x}{3} \left(\frac{k - 76x^2}{3\pi} \right)^{\frac{1}{2}}$$

$$\Rightarrow \frac{45}{19}x^2 = x \left(\frac{k - 76x^2}{3\pi} \right)^{\frac{1}{2}} ; x \neq 0$$

$$\Rightarrow \frac{45}{19}x = \left(\frac{k - 76x^2}{3\pi} \right)^{\frac{1}{2}} \Rightarrow \left(\frac{45}{19} \right)^2 x^2 = \frac{k - 76x^2}{3\pi}$$

$$\Rightarrow \left(\frac{45}{19} \right)^2 x^2 = r^2 \Rightarrow \frac{x^2}{r^2} = \left(\frac{19}{45} \right)^2$$

$$\Rightarrow \frac{x}{r} = \frac{19}{45}$$

7. The area of the region bounded by $y^2 = 8x$ and $y^2 = 16(3 - x)$ is equal to :-


(A) $\frac{32}{3}$ (B) $\frac{40}{3}$ (C) 16 (D) 19

Official Ans. by NTA (C)

Allen Ans. (C)

Sol. $y^2 = 8x ; y^2 = 16(3 - x)$

$$y^2 = -16(x - 3)$$



finding their intersection pts.

$$y^2 = 8x \text{ & } y^2 = -16(x - 3)$$

$$8x = -16x + 48$$

$$24x = 48$$

$$x = 2; y = \pm 4$$

$$A = 2 \int_0^4 (x_R - x_L) dy$$

Required Area

$$\begin{aligned} &= 2 \int_0^4 \left(3 - \frac{y^2}{16} - \frac{y^2}{8} \right) dy \\ &= 2 \left(3y - \frac{y^3}{3 \times 16} - \frac{y^3}{3 \times 8} \right)_0^4 \\ &= 2 \left(3 \times 4 - \frac{4 \times 4 \times 4}{3 \times 16} - \frac{4 \times 4 \times 4 \times 2}{3 \times 8 \times 2} \right) \\ &= 2 \left(12 - \frac{4}{3} - \frac{8}{3} \right) = 2 \times 12 \left(1 - \frac{1}{3} \right) = 2 \times 12 \times \frac{2}{3} = 16 \end{aligned}$$

8. If $\int \frac{1}{x} \sqrt{\frac{1-x}{1+x}} dx = g(x) + c$, $g(1) = 0$, then $g\left(\frac{1}{2}\right)$ is equal to :

(A) $\log_e \left(\frac{\sqrt{3}-1}{\sqrt{3}+1} \right) + \frac{\pi}{3}$ (B) $\log_e \left(\frac{\sqrt{3}+1}{\sqrt{3}-1} \right) + \frac{\pi}{3}$
 (C) $\log_e \left(\frac{\sqrt{3}+1}{\sqrt{3}-1} \right) - \frac{\pi}{3}$ (D) $\frac{1}{2} \log_e \left(\frac{\sqrt{3}-1}{\sqrt{3}+1} \right) - \frac{\pi}{6}$

Official Ans. by NTA (A)

Allen Ans. (A)

Sol. $\int \frac{1}{x} \sqrt{\frac{1-x}{1+x}} dx = g(x) + c$

$$\text{Put } x = \cos 2\theta$$

$$dx = -2\sin 2\theta \cdot d\theta$$

$$= \int \frac{1}{\cos 2\theta} \tan \theta (-4 \sin \theta \cdot \cos \theta) d\theta$$

$$= \int \frac{1}{\cos 2\theta} (-4 \sin^2 \theta) d\theta$$

$$= -\frac{2}{2} \ln |\sec 2\theta + \tan 2\theta| + 2\theta + c$$

$$= \ln |\sec 2\theta - \tan 2\theta| + 2\theta + c$$

$$= \ln \left| \frac{1 - \sqrt{1 - x^2}}{x} \right| + \cos^{-1} x + c$$

$$\underbrace{\left| \frac{1 - \sqrt{1 - x^2}}{x} \right|}_{g(x)} + \cos^{-1} x + c$$

$$\therefore g(1) = 0$$

$$g(x) = \ln \left| \frac{1 - \sqrt{1 - x^2}}{x} \right| + \cos^{-1} x$$

$$g\left(\frac{1}{2}\right) = \ln \left| 2 - \sqrt{3} \right| + \frac{\pi}{3}$$

$$g\left(\frac{1}{2}\right) = \ln \left| \frac{\sqrt{3}-1}{\sqrt{3}+1} \right| + \frac{\pi}{3}$$

9. If $y = y(x)$ is the solution of the differential equation $x \frac{dy}{dx} + 2y = xe^x$, $y(1) = 0$, then the local

maximum value of the function $z(x) = x^2 y(x) - e^x$, $x \in \mathbb{R}$ is :

(A) $1 - e$ (B) 0 (C) $\frac{1}{2}$ (D) $\frac{4}{e} - e$

Official Ans. by NTA (D)

Allen Ans. (D)

$$x \frac{dy}{dx} + 2y = xe^x$$

$$\frac{dy}{dx} + \frac{2y}{x} = e^x$$

$$\text{I.F.} = x^2$$

$$y \cdot x^2 = \int x^2 e^x dx$$

$$\begin{aligned}
 &= \int e^x (x^2 + 2x - 2x - 2 + 2) dx \\
 yx^2 &= e^x (x^2 - 2x + 2) + c \\
 y(1) &= 0 \\
 0 &= e(1 + 0) + c \\
 c &= -e
 \end{aligned}$$

$$\begin{aligned}
 z(x) &= x^2 y(x) - e^x \\
 &= e^x (x^2 - 2x + 2) - e - e^x \\
 &= e^x (x - 1)^2 - e
 \end{aligned}$$

$$\begin{aligned}
 \frac{dz}{dx} &= e^x \cdot 2(x-1) + e^x (x-1)^2 = 0 \\
 x^2(x-1)(2+x-1) &= 0 \\
 e^x(x-1)(x+1) &= 0 \\
 x &= -1, 1
 \end{aligned}$$



$x = -1$ local maxima. Then maximum value is

$$z(-1) = \frac{4}{e} - e$$

10. If the solution of the differential equation

$$\frac{dy}{dx} + e^x (x^2 - 2) y = (x^2 - 2x)(x^2 - 2)e^{2x} \quad \text{satisfies}$$

$y(0) = 0$, then the value of $y(2)$ is _____.

- (A) -1 (B) 1 (C) 0 (D) e

Official Ans. by NTA (C)

Allen Ans. (C)

$$\begin{aligned}
 \text{Sol. } I.F. &= e^{\int e^x (x^2 - 2) dx} = e^{\int e^x (x^2 - 2x + 2x - 2) dx} \\
 &= e^{e^x (x^2 - 2x)}
 \end{aligned}$$

$$y \cdot e^{e^x (x^2 - 2x)} = \int e^{e^x (x^2 - 2x)} e^x (x^2 - 2x)(x^2 - 2) e^x dx$$

Let $e^x (x^2 - 2x) = t$

$$\text{So, } y \cdot e^{e^x (x^2 - 2x)} = \int e^t \cdot t dt$$

At $x = 0, t = 0$

$$x = 2, t = 0$$

$$= t \cdot e^t - e^t + c$$

$$x = 0 ; 0 \cdot 1 = 0 - 1 + c \Rightarrow c = 1$$

$$\text{for } x = 2; y \cdot 1 = 0 - 1 + 1 = 0$$

$$y(2) = 0$$

11. If m is the slope of a common tangent to the curves $\frac{x^2}{16} + \frac{y^2}{9} = 1$ and $x^2 + y^2 = 12$, then $12m^2$ is equal to :
- (A) 6 (B) 9
(C) 10 (D) 12

Official Ans. by NTA (B)

Allen Ans. (B)

$$\begin{aligned}
 \text{Sol. } \frac{x^2}{16} + \frac{y^2}{9} &= 1 \\
 \text{equation of tangent to the ellipse is} \\
 y &= mx \pm \sqrt{a^2 m^2 + b^2} \\
 y &= mx \pm \sqrt{16m^2 + 9} \\
 x^2 + y^2 &= 12
 \end{aligned}$$

equation of tangent to the circle is

$$y = mx \pm \sqrt{12} \sqrt{1+m^2} \quad \dots\text{(ii)}$$

for common tangent equate eq. (i) and (ii)

$$\Rightarrow 16m^2 + 9 = 12(1 + m^2)$$

$$16m^2 - 12m^2 = 3$$

$$4m^2 = 3$$

$$12m^2 = 9$$

12. The locus of the mid point of the line segment joining the point (4, 3) and the points on the ellipse $x^2 + 2y^2 = 4$ is an ellipse with eccentricity :

$$\begin{array}{ll}
 \text{(A) } \frac{\sqrt{3}}{2} & \text{(B) } \frac{1}{2\sqrt{2}} \\
 \text{(C) } \frac{1}{\sqrt{2}} & \text{(D) } \frac{1}{2}
 \end{array}$$

Official Ans. by NTA (C)

Allen Ans. (C)

$$\begin{aligned}
 \text{Sol. } \frac{x^2}{4} + \frac{y^2}{2} &= 1 \\
 P(4,3) \bullet \bullet \bullet Q(2\cos\theta, \sqrt{2}\sin\theta) &
 \end{aligned}$$

Coordinate of D is

$$\left(\frac{2\cos\theta + 4}{2}, \frac{\sqrt{2}\sin\theta + 3}{2} \right) \equiv (h, k)$$

$$\frac{2h - 4}{2} = \cos\theta \quad \dots\text{(i)}$$

$$\frac{2k - 3}{\sqrt{2}} = \sin\theta \quad \dots\text{(ii)}$$

$(i)^2 + (ii)^2$, then we get

$$\left(\frac{2h-4}{2}\right)^2 + \left(\frac{2k-3}{\sqrt{2}}\right)^2 = 1 \Rightarrow \frac{(x-2)^2}{1} + \frac{\left(y-\frac{3}{2}\right)^2}{\left(\frac{1}{2}\right)} = 1$$

\therefore Required eccentricity is

$$e = \sqrt{1 - \frac{1}{2}} = \frac{1}{\sqrt{2}}$$

13. The normal to the hyperbola $\frac{x^2}{a^2} - \frac{y^2}{9} = 1$ at the point $(8, 3\sqrt{3})$ on it passes through the point :
- (A) $(15, -2\sqrt{3})$ (B) $(9, 2\sqrt{3})$
 (C) $(-1, 9\sqrt{3})$ (D) $(-1, 6\sqrt{3})$

Official Ans. by NTA (C)

Allen Ans. (C)

Sol. $\frac{x^2}{a^2} - \frac{y^2}{9} = 1 : (8, 3\sqrt{3})$ lie on Hyperbola then

$$\frac{64}{a^2} - \frac{27}{9} = 1 \Rightarrow a^2 = \frac{64}{4} = 16$$

equation of normal at $(8, 3\sqrt{3})$:

$$\frac{16x}{8} + \frac{9y}{3\sqrt{3}} = 16 + 9$$

$$2x + \sqrt{3}y = 25$$

Check options.

14. If the plane $2x + y - 5z = 0$ is rotated about its line of intersection with the plane $3x - y + 4z - 7 = 0$ by an angle of $\frac{\pi}{2}$, then the plane after the rotation passes through the point :
- (A) $(2, -2, 0)$ (B) $(-2, 2, 0)$
 (C) $(1, 0, 2)$ (D) $(-1, 0, -2)$

Official Ans. by NTA (C)

Allen Ans. (C)

Sol. $(2x + y - 5z) + \lambda(3x - y + 4z - 7) = 0$

Rotated by $\pi/2$

$$(2 + 3\lambda)x + (1 - \lambda)y + (-5 + 4\lambda)z - 7\lambda = 0$$

$$2x + y - 5z = 0$$

$$2(2 + 3\lambda) + (1 - \lambda) - 5(-5 + 4\lambda) = 0$$

$$\Rightarrow 4 + 6\lambda + 1 - \lambda + 25 - 20\lambda = 0$$

$$30 = 15\lambda$$

$$\lambda = 2$$

$$\text{Required plane : } 8x - y + 3z - 14 = 0$$

Check options

15. If the lines $\vec{r} = (\hat{i} - \hat{j} + \hat{k}) + \lambda(3\hat{j} - \hat{k})$ and $\vec{r} = (\alpha\hat{i} - \hat{j}) + \mu(2\hat{i} - 3\hat{k})$ are co-planar, then distance of the plane containing these two lines from the point $(\square, 0, 0)$ is :

$$(A) \frac{2}{9} \quad (B) \frac{2}{11}$$

$$(C) \frac{4}{11} \quad (D) 2$$

Official Ans. by NTA (B)

Allen Ans. (B)

$$\vec{r} = (\hat{i} - \hat{j} + \hat{k}) + \lambda(3\hat{j} - \hat{k}) \dots L_1$$

$$\vec{r} = (\alpha\hat{i} - \hat{j}) + \mu(2\hat{i} - 3\hat{k}) \dots L_2$$

\square L_1 and L_2 are coplanar

$$\therefore \begin{vmatrix} 0 & 3 & -1 \\ 2 & 0 & -3 \\ (1-\alpha) & 0 & 1 \end{vmatrix} = 0$$

$$-3(2 + 3(1 - \square)) = 0$$

$$2 + 3 - 3\square = 0$$

$$\square 3\square = 5$$

$$\Rightarrow \alpha = \frac{5}{3}$$

Now,

$$\vec{n} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 0 & 3 & -1 \\ 2 & 0 & -3 \end{vmatrix} = \hat{i}(-9) - \hat{j}(2) + \hat{k}(-6)$$

$$= (9, 2, 6)$$

Equation of plane :

$$9(x - 1) + 2(y + 1) + 6(z - 1) = 0$$

$$9x + 2y + 6z - 13 = 0$$

Perpendicular distance from $(\square, 0, 0)$

$$= \frac{\left| 9 \cdot \frac{5}{3} + 0 + 0 - 13 \right|}{\sqrt{81 + 36 + 4}} = \frac{2}{\sqrt{121}} = \frac{2}{11}$$

16. Let $\vec{a} = \hat{i} + \hat{j} + 2\hat{k}$, $\vec{b} = 2\hat{i} - 3\hat{j} + \hat{k}$ and $\vec{c} = \hat{i} - \hat{j} + \hat{k}$ be three given vectors. Let \vec{v} be a vector in the plane of \vec{a} and \vec{b} whose projection on \vec{c} is $\frac{2}{\sqrt{3}}$. If $\vec{v} \cdot \hat{j} = 7$, then $\vec{v} \cdot (\hat{i} + \hat{k})$ is equal to :
- (A) 6 (B) 7 (C) 8 (D) 9

Official Ans. by NTA (D)

Allen Ans. (D)

Sol. $\vec{v} = \lambda \vec{a} + \mu \vec{b}$

$$\vec{v} = \lambda(1,1,2) + \mu(2,-3,1)$$

$$\vec{v} = (\lambda + 2\mu, \lambda - 3\mu, 2\lambda + \mu)$$

$$\vec{v} \cdot \hat{j} = 7$$

$$\vec{v} \cdot \frac{\vec{c}}{|\vec{c}|} = \frac{2}{\sqrt{3}}$$

$$\lambda - 3\mu = 7$$

$$\vec{v} \cdot \vec{c} = 2$$

$$\lambda + 2\mu - \lambda + 3\mu + 2\lambda + \mu = 2$$

$$2\lambda + 6\mu = 2$$

$$\lambda + 3\mu = 1$$

$$\lambda - 3\mu = 7$$

$$2\lambda = 8$$

$$\lambda = 4$$

$$\mu = -1$$

We get $\vec{v} = (2, 7, 7)$

17. The mean and standard deviation of 50 observations are 15 and 2 respectively. It was found that one incorrect observation was taken such that the sum of correct and incorrect observations is 70. If the correct mean is 16, then the correct variance is equal to :

(A) 10 (B) 36 (C) 43 (D) 60

Official Ans. by NTA (C)

Allen Ans. (C)

Sol. No. of observations: - 50

$$\text{mean}(\bar{x}) = 15$$

$$\text{Standard deviation } (\sigma) = 2$$

Let incorrect observation is x_1 & correct observation is (x'_1)

$$\text{Given } x_1 + x'_1 = 70$$

$$\bar{x} = \frac{x_1 + x_2 + \dots + x_{50}}{50} = 15 \text{ (given)}$$

$$\Rightarrow x_1 + x_2 + \dots + x_{50} = 750 \quad \dots \text{(i)}$$

Now

Mean of correct observation is 16

$$\frac{x'_1 + x'_2 + \dots + x'_{50}}{50} = 16$$

$$x'_1 + x'_2 + x'_3 + \dots + x'_{50} = 16 \times 50 \quad \dots \text{(ii)}$$

$$\text{eq. (ii)} - \text{eq. (i)}$$

$$\Rightarrow x'_1 - x_1 = 16 \times 50 - 15 \times 50$$

$$x'_1 - x_1 = 50 \text{ & } x_1 + x'_1 = 70$$

$$x'_1 = 60$$

$$x_1 = 10$$

$$\Rightarrow 4 = \frac{x_1^2 + x_2^2 + \dots + x_{50}^2}{50} - 15^2 \quad \dots \text{(iii)}$$

$$\Rightarrow \sigma^2 = \frac{x'_1^2 + x'_2^2 + \dots + x'_{50}^2}{50} - 16^2 \quad \dots \text{(iv)}$$

from (iii)

$$\Rightarrow 4 = \frac{(10)^2}{50} + \frac{x_2^2 + x_3^2 + \dots + x_{50}^2}{50} - 225$$

$$\Rightarrow 4 = 2 - 225 + \frac{(x_2^2 + x_3^2 + \dots + x_{50}^2)}{50}$$

$$\Rightarrow 227 = \frac{(x_2^2 + x_3^2 + \dots + x_{50}^2)}{50}$$

From (iv)

$$\sigma^2 = \frac{(60)^2}{50} + \left(\frac{x_2^2 + x_3^2 + \dots + x_{50}^2}{50} \right) - (16)^2$$

$$\sigma^2 = \frac{60 \times 60}{50} + 227 - 256$$

$$\sigma^2 = 72 + 227 - 256$$

$$\sigma^2 = 43$$

18. $16 \sin(20^\circ) \sin(40^\circ) \sin(80^\circ)$ is equal to :

(A) $\sqrt{3}$ (B) $2\sqrt{3}$ (C) 3 (D) $4\sqrt{3}$

Official Ans. by NTA (B)

Allen Ans. (B)

$$\text{Sol. } 16 \sin 20^\circ \sin 40^\circ \sin 80^\circ$$

$$= 16 \sin 40^\circ \sin 20^\circ \sin 80^\circ$$

$$= 4(4 \sin(60 - 20) \sin(20) \sin(60 + 20))$$

$$= 4 \times \sin(3 \times 20^\circ)$$

$$[\because \sin 3\theta = 4 \sin(60 - \theta) \times \sin \theta \times \sin(60 + \theta)]$$

$$= 4 \times \sin 60^\circ$$

$$= 4 \times \frac{\sqrt{3}}{2} = 2\sqrt{3}$$


19. If the inverse trigonometric functions take principal values, then
 $\cos^{-1}\left(\frac{3}{10}\cos\left(\tan^{-1}\left(\frac{4}{3}\right)\right) + \frac{2}{5}\sin\left(\tan^{-1}\left(\frac{4}{3}\right)\right)\right)$ is equal to :
(A) 0 (B) $\frac{\pi}{4}$ (C) $\frac{\pi}{3}$ (D) $\frac{\pi}{6}$

Official Ans. by NTA (C)

Allen Ans. (C)

Sol. Let

$$\tan^{-1}\frac{4}{3} = \theta \Rightarrow \tan\theta = \frac{4}{3}$$



$$E = \cos^{-1}\left(\frac{3}{10}\cos\theta + \frac{2}{5}\sin\theta\right)$$

$$= \cos^{-1}\left(\frac{3}{10} \times \frac{3}{5} + \frac{2}{5} \cdot \frac{4}{5}\right)$$

$$= \cos^{-1}\left(\frac{9}{50} + \frac{8}{25}\right) = \cos^{-1}\left(\frac{25}{50}\right) = \cos^{-1}\left(\frac{1}{2}\right) = \frac{\pi}{3}$$

20. Let $r \in \{p, q, \sim p, \sim q\}$ be such that the logical statement $r \vee (\sim p) \Rightarrow (p \wedge q) \vee r$ is a tautology. Then 'r' is equal to :
(A) p (B) q (C) $\sim p$ (D) $\sim q$

Official Ans. by NTA (C)

Allen Ans. (C)

Sol. By options

(1)

$p=r$	q	$\sim p$	$r \vee (\sim p)$	$(p \wedge q)$	$(p \wedge q) \vee r$	$r \vee (\sim p)$ \Rightarrow $(p \wedge q) \vee r$
T	F	F	T	F	T	T
F	T	T	T	F	F	F
T	T	F	T	T	T	T
F	F	T	T	F	F	F

(2)

p	$\sim p$	$r \vee (\sim p)$	$q=r$	$(p \wedge q)$	$(p \wedge q) \vee r$	$r \vee (\sim p)$ \Rightarrow $(p \wedge q) \vee r$
T	F	T	T	T	T	T
F	T	T	F	F	T	T
T	F	F	F	F	F	T
F	T	T	F	F	T	F

(3)

p	q	$r = \sim p$	$r \vee (\sim p)$	$(p \wedge q)$	$(p \wedge q) \vee r$	$r \vee (\sim p)$ \Rightarrow $(p \wedge q) \vee r$
T	T	F	F	T	T	T
F	T	T	T	F	T	T
T	F	F	F	F	F	T
F	F	T	T	F	T	T

(4)

$\sim p$	p	q	$r \vee (\sim p)$	$r = \sim q$	$(p \wedge q)$	$(p \wedge q) \vee r$	$r \vee (\sim p)$ \Rightarrow $(p \wedge q) \vee r$
F	T	T	F	F	T	T	T
F	T	F	T	T	F	T	T
T	F	T	T	F	F	F	F
T	F	F	T	T	F	T	T

Now final answer is option no. 3.

SECTION-B

1. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ satisfy $f(x+y) = 2^x f(y) + 4^y f(x), \forall x, y \in \mathbb{R}$. If $f(2) = 3$, then $14 \cdot \frac{f'(4)}{f'(2)}$ is equal to _____.

Official Ans. by NTA (248)

Allen Ans. (248)

Sol. Put $y = 2$

$$f(x+y) = 2^x f(y) + 4^y f(x).$$

$$f(x+2) = 2^x \cdot 3 + 16 f(x)$$

$$f(x+2) = 16 f(x) + 3 \cdot 2^x \ln 2$$

$$f(4) = 16 f(2) + 12 \ln 2 \quad \dots \text{(i)}$$

$$f(y+2) = 4f(y) + 3 \cdot 4^y$$

$$f'(y+2) = 4f'(y) + 3 \cdot 4^y \ln 4$$

$$f(4) = 4f(2) + 96 \ln 2 \quad \dots(ii)$$

solving eq. (i) and (ii), we get

$$f(2) = 7 \ln 2$$

from equation (i), we get

$$f(4) = 124 \ln 2$$

$$\text{Now, } \Rightarrow 14 \cdot \frac{f'(4)}{f'(2)}$$

$$14 \times \frac{124 \ln 2}{7 \ln 2}$$

$$= 248.$$

2. Let p and q be two real numbers such that $p + q = 3$ and $p^4 + q^4 = 369$. Then $\left(\frac{1}{p} + \frac{1}{q}\right)^{-2}$ is equal to _____.

Official Ans. by NTA (4)

Allen Ans. (4)

$$\text{Sol. } p + q = 3 \quad p^4 + q^4 = 369$$

$$\left(\frac{1}{p} + \frac{1}{q}\right)^{-2}$$

$$(p + q)^2 = 9$$

$$p^2 + q^2 = 9 - 2pq$$

$$\frac{1}{\left(\frac{1}{p} + \frac{1}{q}\right)^2} = \frac{(pq)^2}{(q+p)^2} = \frac{(pq)^2}{9}$$

$$p^4 + q^4 = (p^2 + q^2)^2 - 2p^2q^2$$

$$369 = (9 - 2pq)^2 - 2(pq)^2$$

$$369 = 81 + 4p^2q^2 - 36pq - 2p^2q^2$$

$$288 = 2p^2q^2 - 36pq$$

$$144 = p^2q^2 - 18pq$$

$$(pq)^2 - 2 \times 9 \times pq + 9^2 = 144 + 9^2$$

$$(pq - 9)^2 = 225$$

$$pq - 9 = \pm 15$$

$$pq = \pm 15 + 9$$

$$pq = 24, -6$$

(24 is rejected because $p^2 + q^2 = 9 - 2pq$ is negative)

$$\frac{(pq)^2}{9} = \frac{1(-6)^2}{9} = 4$$

3. If $z^2 + z + 1 = 0$, $z \in \mathbb{C}$, then $\left| \sum_{n=1}^{15} \left(Z^n + (-1)^n \frac{1}{Z^n} \right)^2 \right|$ is equal to _____.

Official Ans. by NTA (2)

Allen Ans. (2)

$$\text{Sol. } z^2 + z + 1 = 0 \Rightarrow z = w, w^2$$

$$\left| \sum_{n=1}^{15} \left(Z^n + (-1)^n \frac{1}{Z^n} \right)^2 \right| = \left| \sum_{n=1}^{15} \left(Z^{2n} + \frac{1}{Z^{2n}} + 2(-1)^n \right) \right|$$

$$= \left| \sum_{n=1}^{15} w^{2n} + \frac{1}{w^{2n}} + 2(-1)^n \right|$$

$$= \left| \frac{w^2(1-w^{30})}{1-w^2} + \frac{\frac{1}{w^2}(1-\frac{1}{w^{30}})}{1-\frac{1}{w^2}} + 2(-1) \right|$$

$$= \left| \frac{w^2(1-1)}{1-w^2} + \frac{\frac{1}{w^2}(1-1)}{1-\frac{1}{w^2}} - 2 \right|$$

$$= |0 + 0 - 2| = 2$$

4. Let $X = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$, $Y = \alpha I + \beta X + \gamma X^2$ and

$$Z = \alpha^2 I - \alpha\beta X + (\beta^2 - \alpha\gamma)X^2, \alpha, \beta, \gamma \in \mathbb{R}$$

$$\text{If } Y^{-1} = \begin{bmatrix} \frac{1}{5} & -\frac{2}{5} & \frac{1}{5} \\ 0 & \frac{1}{5} & -\frac{2}{5} \\ 0 & 0 & \frac{1}{5} \end{bmatrix}, \text{ then } (\alpha - \beta + \gamma)^2 \text{ is equal to } \text{_____}.$$

Official Ans. by NTA (100)

Allen Ans. (100)

$$X = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}, X^2 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$Y = \begin{bmatrix} \alpha & \beta & \gamma \\ 0 & \alpha & \beta \\ 0 & 0 & \alpha \end{bmatrix}, Z = \begin{bmatrix} \alpha^2 & -\alpha\beta & \beta^2 - \alpha\gamma \\ 0 & \alpha^2 & -\alpha\beta \\ 0 & 0 & \alpha^2 \end{bmatrix}$$

$$Y \cdot Y^{-1} = I$$

$$\begin{bmatrix} \alpha & \beta & \gamma \\ 0 & \alpha & \beta \\ 0 & 0 & \alpha \end{bmatrix} \begin{bmatrix} \frac{1}{5} & -\frac{2}{5} & \frac{1}{5} \\ 0 & \frac{1}{5} & -\frac{2}{5} \\ 0 & 0 & \frac{1}{5} \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\frac{\alpha}{5} = 1 \Rightarrow \alpha = 5$$

$$-\frac{2}{5}\alpha + \frac{\beta}{5} = 0 \Rightarrow \beta = 10$$

$$\frac{\alpha}{5} - \frac{2\beta}{5} + \frac{\gamma}{5} = 0 \Rightarrow \gamma = 15$$

$$\Rightarrow (\alpha - \beta + \gamma)^2 = (5 - 10 + 15)^2 = 100$$

5. The total number of 3-digit numbers, whose greatest common divisor with 36 is 2, is _____. .

Official Ans. by NTA (150)

Allen Ans. (150)

Sol. $36 = 2 \times 2 \times 3 \times 3$

Number should be odd multiple of 2 and does not having factor 3 and 9

Odd multiple of 2 are

102, 106, 110, 114 998 (225 no.)

No. of multiples of 3 are

102, 114, 126 990 (75 no.)

Which are also included multiple of 9

Hence,

$$\text{Required} = 225 - 75 = 150$$

6. If $\binom{40}{0} + \binom{41}{1} + \binom{42}{2} + \dots + \binom{60}{20} = \frac{m}{n} \binom{60}{20}$, m

and n are coprime, then m + n is equal to _____. .

Official Ans. by NTA (102)

Allen Ans. (102)

Sol. ${}^{40}C_0 + {}^{41}C_1 + {}^{42}C_2 + \dots + {}^{59}C_{19} + {}^{60}C_{20}$

$$\left(\frac{1}{41} + 1\right) {}^{41}C_1 + {}^{42}C_2 + \dots$$

$$\left[\frac{42}{41} \left(\frac{2}{42}\right) + 1\right] {}^{42}C_2 + {}^{43}C_3 + \dots$$

$$\left(\frac{2}{41} + 1\right) {}^{42}C_2 + {}^{43}C_3 + \dots$$

$$\left(\frac{43}{41} \times \frac{3}{43} + 1\right) {}^{43}C_3 + {}^{44}C_4 + \dots$$

$$\frac{3+41}{41} \cdot {}^{43}C_3 + \dots$$

Similarly :

$$\frac{20+41}{41}$$

$$\Rightarrow m = 61 ; n = 41$$

$$m + n = 102$$

7. If $a_1 (> 0), a_2, a_3, a_4, a_5$ are in a G.P., $a_2 + a_4 = 2a_3 + 1$ and $3a_2 + a_3 = 2a_4$, then $a_2 + a_4 + 2a_5$ is equal to _____. .

Official Ans. by NTA (40)

Allen Ans. (40)

Sol. $a_1 > 0, a_2, a_3, a_4, a_5 \rightarrow \text{G.P.}$

$$3a_2 + a_3 = 2a_4$$

$$3ar + ar^2 = 2ar^3$$

$$3 + r = 2r^2$$

$$2r^2 - r - 3 = 0$$

$$r = -1 \text{ & } r = \frac{3}{2}$$

$$a_2 + a_4 = 2a_3 + 1$$

$$ar + ar^3 = 2ar^2 + 1$$

$$a(r + r^3 - 2r^2) = 1$$

$$a \left(\frac{3}{2} + \frac{27}{8} - \frac{18}{4} \right) = 1$$

$$a = \frac{8}{3}$$

$$\text{When } r = -1, a = -\frac{1}{4} \text{ (rejected, } a_1 > 0\text{)}$$

$$r = \frac{2}{3}, a = \frac{8}{3} \text{ (selected)}$$

Now

$$a_2 + a_4 + 2a_5$$

$$= \frac{8}{3} \times \frac{3}{2} + \frac{8}{3} \times \frac{27}{8} + 2 \times \frac{8}{3} \times \frac{81}{16}$$

$$= 4 + 9 + 27 = 40$$

8. The integral $\frac{24}{\pi} \int_0^{\sqrt{2}} \frac{(2-x^2)}{(2+x^2)\sqrt{4+x^4}} dx$ is equal to _____. .

Official Ans. by NTA (3)

Allen Ans. (3)

Sol. $\frac{24}{\pi} \int_0^{\sqrt{2}} \frac{(2-x^2)}{(x^2+2)\sqrt{4+x^4}} dx$

$$\frac{24\sqrt{2}}{\pi} \int_0^{\frac{\pi}{2}} \frac{x^2 \left(\frac{2}{x^2} - 1 \right) dx}{x \left(x + \frac{2}{x} \right) \times x \sqrt{\frac{4}{x^2} + x^2}}$$

$$\frac{24\sqrt{2}}{\pi} \int_0^{\frac{\pi}{2}} \frac{\left(\frac{2}{x^2} - 1 \right) dx}{\left(x + \frac{2}{x} \right) \sqrt{\left(x + \frac{2}{x} \right)^2 - 4}}$$

$$x + \frac{2}{x} = t$$

$$dt = \left(1 - \frac{2}{x^2} \right) dx$$

$$I = -\frac{24}{\pi} \int \frac{dt}{t \sqrt{t^2 - 4}}$$

$$= -\frac{24}{\pi} \times \frac{1}{2} \sec^{-1} \left(\frac{x + \frac{2}{x}}{2} \right) \Big|_0^{\frac{\pi}{2}}$$

$$= -\frac{12}{\pi} \left[\sec^{-1} \left(\frac{2\sqrt{2}}{2} \right) - \sec^{-1} (\infty) \right]$$


$$= -\frac{12}{\pi} \left[\frac{\pi}{4} - \frac{2\pi}{2 \times 2} \right] = -\frac{12}{\pi} \left[-\frac{\pi}{4} \right]$$

$$= 3$$

9. Let a line L_1 be tangent to the hyperbola $\frac{x^2}{16} - \frac{y^2}{4} = 1$ and let L_2 be the line passing through the origin and perpendicular to L_1 . If the locus of the point of intersection of L_1 and L_2 is $(x^2 + y^2)^2 = \alpha x^2 + \beta y^2$, then $\alpha + \beta$ is equal to _____.

Official Ans. by NTA (12)

Allen Ans. (12)



$$m_1 = \frac{\sec \theta \times 2}{4(\tan \theta)} = \frac{\sec \theta}{2 \tan \theta}$$

$$m_2 = \frac{k}{h}$$

$$m_1 m_2 = -1$$

$$\frac{k \cdot \sec \theta}{h \cdot 2 \tan \theta} = -1$$

$$\frac{k}{2h \sin \theta} = -1$$

$$\sin \theta = \frac{-k}{2h} \quad \cos \theta = \frac{\sqrt{4h^2 - k^2}}{2h}$$

also

$$\frac{h \sec \theta}{4} - \frac{k \tan \theta}{2} = 1$$

$$\frac{h}{4} \frac{2h}{\sqrt{4h^2 - k^2}} - \frac{k}{2} \left(\frac{-k}{\sqrt{4h^2 - k^2}} \right) = 1$$

$$h^2 + k^2 = 2\sqrt{4h^2 - k^2}$$

$$(x^2 + y^2)^2 = 4(4x^2 - y^2)$$

$$(x^2 + y^2)^2 = 16x^2 - 4y^2$$

$$\alpha = 16, \beta = -4$$

$$\alpha + \beta = 16 - 4 = 12$$

10. If the probability that a randomly chosen 6-digit number formed by using digits 1 and 8 only is a multiple of 21 is p , then $96p$ is equal to _____.

Official Ans. by NTA (33)

Allen Ans. (33)

Sol. $2 \times 2 \times 2 \times 2 \times 2 \times 2 = 64$

Divisible by 21 when divided by 3.

Case – I : All 1 \rightarrow (1)

Case – II : All 8 \rightarrow (1)

Case – III : 3 ones & 3 eights

$$\frac{6!}{3! \times 3!} = 20$$

$$\text{Required probability } \therefore p = \frac{22}{64}$$

$$96p = 96 \times \frac{22}{64} = 33$$